

Econ-240A (1st half - Fall 2008)
Note on Completeness 2008-10-19 (v2)
Owen Ozier

The idea, expressed in CB Theorem 6.2.25, that a particular sufficient statistic is complete only if (among other conditions from the exponential family representation) $\{(w_1(\theta), \dots, w_k(\theta)) : \theta \in \Theta\}$ contains an open set in \mathbb{R}^k , has gone more or less without explanation; let me try to give an intuitive one based on some examples.

Suppose that we have random variables drawn from the $\text{Exponential}(\theta)$ distribution (in the sense of CB equation 3.3.11). It is straightforward to see that this family of distributions is an exponential family (the CB equation 3.4.1 meaning, now):

$$\frac{1}{\theta} e^{-x/\theta} \mathbb{1}(x \geq 0) = h(x)c(\theta) \exp\left(\sum_{i=1}^k w_i(\theta)t_i(x)\right) \text{ where: } \begin{array}{l} h(x) = \mathbb{1}(x \geq 0) \\ c(\theta) = \frac{1}{\theta} \\ k = 1 \\ w_1(\theta) = \frac{1}{\theta} \\ t_1(x) = x \end{array}$$

Ordinarily, we consider the parameter space $\theta \in \Theta = (0, \infty)$. Since $w_1(\theta) = 1/\theta$ is a continuous function, then because Θ contains an open set in $\mathbb{R}^k = \mathbb{R}^1 = \mathbb{R}$ (in fact, it is one), so does $\{w_1(\theta) : \theta \in \Theta\}$. Thus, by Theorem 6.2.25, the statistic $T(\mathbf{X}) = \sum_{i=1}^n t_1(X_i) = \sum_{i=1}^n X_i$ is complete (and by Theorem 6.2.10, sufficient).

What if Θ didn't include an open set (and thus $w(\Theta)$ also didn't)?

Suppose $\Theta = \{1, 2\}$. Then in this case, we may be able to construct a function $g(\cdot)$, based only on $T(\mathbf{X}) = \sum_{i=1}^n X_i$, that shows that T is not complete. Referring back to the definition of completeness (6.2.21), paraphrased here:

$$\text{If } E[g(T)] = 0 \forall \theta \implies P(g(T) = 0) = 1 \forall \theta, \text{ then } T(X) \text{ is a complete statistic.}$$

Note that $E[T] = n\theta$ and $E[T^2] = n^2\theta^2 + n\theta^2$ (you can work out why this is true). Thus, $E[T^2/(n^2+n)] = \theta^2$, and $E[T/n] = \theta$.

To construct a $g(\cdot)$ that shows that T is not complete, we need $E[g(T)] = 0 \forall \theta$, but now, we know that θ could only take on two values: 1 and 2, so we only need an expression that is zero for those two values. If we had a function whose expectation were $(\theta - 1)(\theta - 2)$, it would satisfy this condition. Multiplying out, we get: $(\theta - 1)(\theta - 2) = \theta^2 - 3\theta + 2$. Based on the notes above, it is easy to construct a function with this expectation:

$$E\left[g(T) = \frac{T^2}{(n^2+n)} - \frac{3T}{n} + 2\right] = \theta^2 - 3\theta + 2 = 0 \forall \theta \in \Theta = \{1, 2\}$$

However, $P[g(T) = 0] \neq 1$ (easy to verify by example), so in this case, $T(\mathbf{X}) = \sum_{i=1}^n X_i$ is not complete.

Why the k -dimensional open set in \mathbb{R}^k ?

The discussion below Theorem 6.2.25 in the textbook discusses a case of this, the set of normals with parameter values $\mathcal{N}(\theta, \theta^2)$. The problem this introduces is that the set of $\{w_i(\theta)\}$ now only contains values on a parabola in \mathbb{R}^2 , so it does not contain any two-dimensional open neighborhoods. Practically, the reason this makes the usual statistics (\bar{X}, S^2) no longer complete, is that it gives us two ways to create functions with expectations related to theta, based on either \bar{X} or S^2 . These two methods may have the same expectation, but different realized values in any given sample. As above, you can construct functions of \bar{X} and S^2 that have expectation zero, but that are not zero with probability 1, so the statistics are no longer complete.

Summary:

In essence, if $\{(w_1(\theta), \dots, w_k(\theta)) : \theta \in \Theta\}$ does not contain an open set in \mathbb{R}^k (it contained only singletons in the first example, and was of too low a dimension in the second example), then it will be possible to construct a function $g(T(\mathbf{X}))$ that has expectation zero but is not always zero. This would, in turn, make it possible to make many different unbiased estimators of θ based only on $T(\mathbf{X})$, so we could not be sure, without further work, whether any of them were actually the UMVU estimator of θ .